



## U-Pb DETRITAL ZIRCON DATA FROM THE PALEOZOIC SIERRA GRANDE FORMATION, NORTH PATAGONIAN MASSIF, ARGENTINA

Uriz, N. J.<sup>(1)</sup>, Cingolani, C. A.<sup>(1, 2)</sup>, Chemale, Jr. F. F.<sup>(3)</sup> and Armstrong, R. A.<sup>(4)</sup>

- (1) División Científica de Geología-Facultad Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paso del Bosque s/n, (1900)-LA PLATA, ARGENTINA. [nuriz@fcnym.unlp.edu.ar](mailto:nuriz@fcnym.unlp.edu.ar)
- (2) CIG (UNLP-CONICET): Centro de Investigaciones Geológicas, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, 1-644, (1900)-LA PLATA, ARGENTINA.
- (3) Laboratorio de Geología Isotópica, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, Porto Alegre, Brazil.
- (4) The Australian National University, Canberra 0200, ACT, Australia.

**Key words:** U-Pb zircon ages, Sierra Grande Formation, North Patagonian Massif, Silurian – Devonian, provenance

### ABSTRACT

The Sierra Grande Formation is a siliciclastic sedimentary unit with oolitic iron levels of economical importance which crops out in the Argentinean North Patagonian Massif. It was deposited within an open marine environment, with littoral to sub-littoral conditions. A Silurian-Lower Devonian age is assigned based on fossil content. It is mainly composed of medium to fine quartz arenites, wackes, silts, shales and conglomerate beds. In order to establish the source rocks for the Silurian-Devonian basin and also to constrain the sedimentation age, different U-Pb geochronological studies have been carried out on detrital zircons. The zircons have been studied with comparative morphology analysis using SEM images to assess the different population types. For dating using U-Pb isotopes in zircon detrital grains, the high resolution SHRIMP and LA-ICP-MS methodologies were used. The predominant obtained age groups are: Neoproterozoic (547–991 Ma), Cambrian (497–546 Ma) and Ordovician (443–495 Ma). All the analyzed samples also comprise Mesoproterozoic ages of about 1009 to 1382 Ma and smaller proportion Paleoproterozoic (1641 to 2248 Ma) and Neo-Archean (2649–2657 Ma) ages. It is important to note that Silurian ages (428 and 440 Ma) were obtained from a group of zircon crystals, constraining the sedimentation age of the Sierra Grande basin to be not older than Middle Silurian. With these isotopic data it is possible to discuss the provenance of the Silurian-Devonian siliciclastic sedimentary cover of the North Patagonian Massif.

### INTRODUCTION

The Sierra Grande Formation (Müller, 1965) is a siliciclastic sedimentary unit with oolitic iron levels of economical importance (Zanettini, 1981 and references therein). This unit shows different exposures in the eastern sector of the Argentinean North Patagonian Massif, being the most representative, the one located close to the town of Sierra Grande in the Río Negro Province (Fig. 1). The southernmost exposures of this unit are located close to the Gastre Town, to the northeast of the Chubut Province. During the last decades, several contributions were published on the fields of tectonic and paleogeography (Ramos, 1984; 1988; Dalla Salda *et al.* 1992), gravimetry (Kostadinoff and Gelós, 1994), paleomagnetism (Rapalini, 1998), sedimentology (Spalletti *et al.* 1991; Spalletti, 1993), petrology and geochronology of basement, sedimentary rocks and intrusive bodies of the North Patagonian Massif (Varela *et al.*, 1991; 1997). Detailed structural analysis carried out by Japas (2001) and especially zircon data on Sierra Grande Formation among others by Pankhurst *et al.* (2006) promoted deeper understanding of the geological processes that have taken place in the region.

The aim of this contribution is to present new U-Pb detrital zircon ages, in order to constrain the sedimentation age of the Sierra Grande sequence and also to get more information regarding the probable source areas that contributed to the basin during the Silurian and Lower Devonian. The data here presented are part of N. Uriz PhD thesis carried out at the Faculty of Natural Sciences, University of La Plata, Argentina.

## GEOLOGICAL FRAMEWORK

The pre-Silurian basement in this sector of the North Patagonian Massif is characterized by several igneous and metamorphic rocks (Giacosa, 1999; Pankhurst *et al.* 2006). The marine siliciclastic sedimentary rocks of the Sierra Grande Formation crops over an erosive unconformity following a patchy trend. It is mainly composed of medium to fine quartz arenites, wackes, silts, shales and conglomerate beds with iron beds. This unit is approximately 2130m thick based on drilling data (Zanettini, 1981).

The Sierra Grande Formation was deposited within an open marine environment, with littoral to sublittoral conditions (de Alba, 1964; Müller, 1965; Nuñez *et al.*, 1975; Cortés, 1981), in wave and storm shallow marine platform areas (Spalletti *et al.*, 1991). A Silurian-Lower Devonian age is assigned based on fossil content (Manceñido and Damborenea, 1984 and references therein). It is important to mention that the first work on heavy minerals from Sierra Grande Formation was presented by Huber-Grünberg (1990).




Figure 1: Regional geological sketch map and sampling localities of the Sierra Grande Formation.

## METHODOLOGY

New U-Pb geochronological studies had been carried out on detrital zircons of the Sierra Grande Fm. The



sample localities are shown in Figure 1. The obtained detrital zircons had been studied with comparative morphology analysis using SEM (scanning electron microscope) to assess the different populations and types. The isotopic ages were obtained by U-Pb LA-ICP-MS (Laboratorio de Geología Isotópica, UFRGS, Porto Alegre, Brazil) and SHRIMP (Research School of Earth Sciences, The Australian National University, Canberra, Australia) methods.

## MORPHOLOGY OF ZIRCONS

To understand the probable source areas of the Silurian-Devonian siliciclastic rocks of the Sierra Grande Formation, new morphological studies on zircon typology were carried out (Huber-Grünberg, 1990). They show that it is possible to discriminate four different groups of populations (A, B, C and D) following the scheme of Pupin (1980):

Group A. It comprises the euhedral to subhedral prismatic and elongated zircons, typical from both volcanic and plutonic sources. Rounded to subrounded subhedral morphologies are the most common and suggest transport before deposition, but still keeping the characteristic long prismatic morphology. These crystals show different degree of internal zonation, related to diverse growing phases, allowing the grouping of them. They show crystal elongation ratios between 2.4 and 4.28, with sizes ranging from 115µm long and 46µm wide, to a maximum of 333µm long and 73µm wide. These types belong to the subgroup of Pupin (1980) known as: P2, P3, P4, P5, R2, R3 and R4 (Figure 2-A).

Group B. Short prismatic crystals, poly-faceted, a few of them have preserved the euhedral morphology. The predominant morphology is the subhedral, with different roundness. The complex zonation patterns and crystalline structures are related to metamorphic sources. They show crystal elongation ratios between 1.8 and 2.7 with an average long/wide ratio of 2.2. The crystal size ranges from 107µm long and 53µm wide, to a maximum of 242µm long and 102µm wide. These types belong to the subgroup of Pupin (1980) known as: S2, S7, S12, S18 and S19 (Figure 2-B).

Group C. It comprises the equidimensional crystals (short prisms), poly-faceted, showing different degree of roundness, within the age range of the Middle Precambrian. They suggest a metamorphic source rock which has been providing the basin with these materials with their characteristically morphologies (Figure 2-C).

Group D. Comprises rounded and equidimensional zircons. Although some of them show internal zonation they can not be discriminated based on their morphologies. They have an age range from the Neo-Archean to the Neoproterozoic (Figure 2-D). In some of the crystals the old cores show overgrowing that suggests that they have suffered other events before their final deposition. On the other hand it is clear the relationship between the degree of roundness and the age, since the more rounded zircon crystals are the Neo-Archean to Neoproterozoic ones, whereas towards younger ages the crystals are sub-rounded.

## RESULTS

For age dating using U-Pb method on zircon detrital grains, the high resolution SHRIMP methodology was used for samples SGS008 and SGLM013, whereas the LA-ICP-MS for samples SGS007, SGLM013 and SGPS017 (Figure 2). The results show a striking similarity regarding the ages obtained using both age-dating methodologies, underlying the importance of the Neoproterozoic materials dated between 547 and 991 Ma.

The age groups are: Neoproterozoic (547–991 Ma), Cambrian between 497 and 546 Ma and Ordovician between 443 and 495 Ma. In a restricted aspect, but present in all the analyzed samples, we register Mesoproterozoic ages close to 1009 to 1382 Ma and in a smaller fashion Paleoproterozoic ages (1641 to 2248 Ma) and Neo-Archean (2649 to 2657 Ma).

It is important to stress the fact that we have got two Silurian ages (428 and 440 Ma) from a group of zircon crystals, constraining the sedimentation age of the Sierra Grande basin to a maximum the Middle Silurian.

The Cambrian ages can be subdivided into two groups, one with ages from 521 to 546 Ma and another one with ages ranging from 501 to 520 Ma (Lower and Middle Cambrian respectively). The Ordovician ages are concentrated in two main ranges, one between 475 and 495 Ma, and the other one between 462 and 472 Ma (Lower and Middle Ordovician respectively).



The Neoproterozoic age is the most representative element of the analyzed samples. The ages show dispersion, comprising Early, Middle and Late Neoproterozoic zircons. A similar trend is displayed for the Mesoproterozoic zircon ages. The recorded ages are coherent for all the samples, since almost all of them show a Late Mesoproterozoic age. The Paleoproterozoic and Neo-Archean ages, although sparse, are present in most of the samples.




Figure 2: U-Pb concordia diagrams and age histograms from the analyzed samples. A, B, C and D: SEM images of the study zircon populations.

## DISCUSSION AND CONCLUSIONS

Using the U-Pb isotopic data on detrital zircons and the respective morphological studies, it is possible to discuss the provenance of the siliciclastic sedimentary rocks of the Silurian-Devonian basin from the north eastern sector of the Argentinean North Patagonian Massif and also to adjust the sedimentation age according to the following scheme:

1-The data reveal a strong influence of Mesoproterozoic-Neoproterozoic, Cambrian and Ordovician sources.

2-These ages can be mainly related to the igneous-metamorphic basement of the North Patagonian Massif, outcropping from the pre-Andean sector towards the Atlantic Ocean (Pankhurst *et al.* 2006). In this basement different lithologies and metamorphic degrees had been recognized, with ages that go from the Late Proterozoic to the Early Paleozoic. These data are coherent with the present contribution. Within the igneous-metamorphic basement the Mina Gonzalito Complex can be mentioned (Caminos and Llambías, 1984; Giacosa, 1987; Linares *et al.*, 1990; Caminos *et al.*, 1994; Varela *et al.*, 1998); the Yaminué Complex (Caminos, 1983; Varela *et al.*, 2001; Basei *et al.*, 2002; Llambías *et al.*, 2002); the Las Piedras Metamórficas Complex (Tickyj *et al.*, 2002), among others, that together are potential source areas for the analysed material.

Locally, the Cambrian-Ordovician ages are represented by the low metamorphic rocks of the El Jaguelito and equivalent units in the area such as the Nahuel Niyeu Formation at the Valcheta region (Caminos and Llambías, 1984; Giacosa, 1987; 1994; 1999; Chernicoff and Caminos, 1996 a, b). Regarding the plutonic intrusive rocks the Punta Sierra Complex can be mentioned, which comprises the Punta Sierra Granite and the Arroyo Salado Granodiorite, with ages of  $476 \pm 4$  Ma (Varela *et al.*, 1998) to  $475 \pm 6$  Ma (Pankhurst *et al.*,



2006).

3- The zircon grains that show plutonic morphologies, with ages ranging from the Late Neoproterozoic to Early Cambrian can be related to those intrusive rocks that preceded the collisional event of the Pampia Terrane (Rapela *et al.*, 1998). On the other hand, there are A-type granite rocks close to the fold and thrust belt of Sierra de la Ventana, that have been linked to the final plutonic event that took place during the postorogenic collisional event (Gregori *et al.*, 2005). This ended with the rhyolites of La Ermita ( $509 \pm 5.3$  Ma), related to the postcollisional tensional event in this sector of Gondwana (Pankhurst *et al.*, 2006). Within the Chadileuvú Block equivalent units to the North Patagonian Massif basement rocks are present (Tickyj *et al.*, 1999a) which can also be interpreted as a source area of detritus.

4-The highest ages of the detrital zircons are Paleoproterozoic and Neo-Archean, that although rare they suggest recycling from old cratons (Río de la Plata, Kalahari?) with a restricted input to the basin.

5-It is important to underline that the Silurian ages (428 – 440 Ma) from two zircon samples allowed us to establish a maximum sedimentation age between the Lower and Middle Silurian for the Sierra Grande Basin.

**ACKNOWLEDGMENTS:** Field and laboratory works were partially supported by Argentine Institutions: CONICET (PIP 5027) and ANPCyT (PICT 07-10829). We are grateful to Marcelo Manassero and Paulina Abre whose reviews of the first English version and comments improved the paper. We are recognized to the Centro de Investigaciones Geológicas (UNLP-CONICET) for their laboratory and equipment facilities. We specially acknowledge to Prof. Moacir Macambira (Belem, Brazil) for his relevant helpful for laboratory work and review the manuscript. We thank to Prof. F. Hervé for his fruitful comments as a reviewer.

## REFERENCES

- Basei, M.A.S., Varela, R., Sato, A.M., Siga Jr., O. and Llambías, E.J., 2002. Geocronología sobre rocas del Complejo Yaminué, Macizo Norpatagónico, Río Negro, Argentina. *Actas 15° Congreso Geológico Argentino* (El Calafate), 3: 117-122, Buenos Aires.
- Camino, R., 1983. Descripción Geológica de las Hojas 39g, Cerro Tapilque y 39h, Chipauquil, provincia de Río Negro. Servicio Geológico Nacional (inédito). Buenos Aires.
- Camino, R. and Llambías, E.J., 1984. El Basamento Cristalino. *Relatorio 9° Congreso Geológico Argentino* (San Carlos de Bariloche), 1: 37-63, Buenos Aires.
- Camino, R., Chernicoff, C. and Varela, R., 1994. Evolucion tectónico-metamórfica y edad del Complejo Yaminué, Basamento pre-andino norpatagónico, República Argentina. 7° Congreso Geológico Chileno (Universidad de Concepción), *Actas II*, 1301–1305.
- Chernicoff, C. J. and Camino, R., 1996a. Estructura y metamorfismo del Complejo Yaminué, Macizo Nordpatagónico oriental, provincia de Río Negro. *Revista de la Asociación Geológica Argentina* 51(2):107-118.
- Chernicoff, C. J. and Camino, R., 1996 b. Estructura y relaciones estratigráficas de la Formación Nahuel Niyeu, Macizo Nordpatagónico oriental, Provincia de Río Negro. *Revista de la Asociación Geológica Argentina* 51(3): 201-212.
- Cortés, J. M. (1981): Estratigrafía cenozoica y estructura al oeste de la Península de Valdés, Chubut. *Consideraciones tectónicas y paleogeográficas. Revista de la Asociación Geológica Argentina*, 36 (4): 424-445.
- Dalla Salda, L.H., Dalziel, I.W.D., Cingolani, C.A. and Varela, R., 1992. Did Taconic Appalachians continue into southern South America? *Geology* 20, 1059–1062.
- De Alba, E., 1964. Descripción geológica de la Hoja 41j "Sierra Grande", provincia de Río Negro. *Carta Geológica y Económica de la República Argentina, Boletín 97, Dirección Nacional de Geología y Minería*
- Giacosa, R., 1987. Caracterización de un sector del basamento metamórfico-migmatítico en el extremo suroriental del Macizo Norpatagónico, provincia de Río Negro. *Actas del X Congreso Geológico Argentino*, 3: 51-54. Tucumán.
- Giacosa, R.E., 1994. El basamento Precámbrico del sector oriental del Macizo Nordpatagónico, Argentina. *Zentralblatt für Geologie und Paläontologie, Teil 1* 1993 (1/2), 89–100.
- Giacosa, R.E., 1999. El basamento pre-Silúrico del extremo Este del Macizo Nordpatagónico y del Macizo del Deseado. *Instituto de Geología y Recursos Minerales, Anales* 29 (5), 118–123.
- Gregori D.A., López, V.L. and Grecco, L.E. 2005. A Late Proterozoic–Early Paleozoic magmatic cycle in Sierra de la Ventana, Argentina. *Journal of South American Earth Sciences*. 19: 155–171.
- Huber-Grünberg, A. 1990. Sedimentologie, Fazies und Herkunft der kambrisch/ordovizischen und silurisch/unterdevonischen Einheiten von Sierra Grande, Patagonien. *Dissertation University of Munich*, 196 pp.
- Japas, M S., 2001. Modelo cinemático neopaleozoico para el sector nororiental del Macizo Norpatagónico, Argentina.



- Journal of Iberian Geology, 27, 91-121
- Kostadinoff, J. and Gelós, E. M., 1994. Análisis de las mediciones gravi-magnetométricas realizadas entre El Fuerte y Arroyo Verde, provincia de Río Negro. *Revista de la Asociación Geológica Argentina*, 49 (1/2): 19-25.
- Linares, E., Ostera, H.A. and Parica, C.A., 1990. Edades radimétricas preliminares del basamento cristalino de las vecindades de Mina Gonzalito y de Valcheta, Provincia de Río Negro, República Argentina. 11° Congreso Geológico Argentino, Actas II, 251–253.
- Llambías, E.J., Varela, R., Basei, M.A.S. and Sato, A.M., 2002. Deformación dúctil y metamorfismo neopaleozoico en el área de Yaminué y su relación con la Fase Orogénica San Rafael. *Actas 15° Congreso Geológico Argentino (El Calafate)*, 3: 117-122, Buenos Aires.
- Manceñido, M. and Damborenea, S., 1984. Megafauna de invertebrados paleozoicos y mesozoicos. In: Ramos, V. (Ed.), *Geología y Recursos Naturales de la Provincia de Río Negro: IX. Congreso Geológico Argentino, Bariloche, Relatorio. Asociación Geológica Argentina, Buenos Aires*, pp. 413–466.
- Müller, H., 1965. Zur altersfrage der Eisenerzlagerstätte Sierra Grande / Río Negro in Nordpatagonien aufgrund neuer fossillunde. *Geologische Rundschau*, 54 (2): 715-732.
- Núñez, E., Bachmann, E. W., Ravazzoli, I., Britos, A., Franchini, M., Lizuaín, A., Sepúlveda, E., 1975. Rasgos geológicos del sector oriental del Macizo de Somuncurá, provincia de Río Negro, República Argentina. II Congreso Iberoamericano de Geología Económica, 4: 247-266.
- Pankhurst, R.J., Rapela, C.W. and Fanning, C.M., 2001. The Mina Gonzalito gneiss: Early Ordovician metamorphism in northern Patagonia. 3° South American Symposium on Isotope Geology, (CD-Rom) Pucón, Chile.
- Pankhurst, R.J., Rapela C.W., Fanning C.M. and Márquez M., 2006. Gondwanide continental collision and the origin of Patagonia. *Earth-Science Reviews* 76: 235–257
- Pupin, J. P., 1980. Zircon and granite petrology. *Contributions to Mineralogy and Petrology*, 73: 207-220.
- Ramos, V. A., 1984. Patagonia: Un continente Paleozoico a la deriva? *Actas del IX Congreso Geológico Argentino*, 2: 311-325. Bariloche.
- Ramos, V.A., 1988. Late Proterozoic–Early Paleozoic of South America - a collisional history. *Episodes* 11 (3), 168–174.
- Rapalini, A., 1998. Syntectonic magnetization of the Mid-Paleozoic Sierra Grande Formation: Further constraints on the tectonic evolution of Patagonia. *Journal of the Geological Society, London*, 155:105-114.
- Rapela, C. W., Pankhurst, R. J., Casquet, C., Baldo, E., Saavedra, J., Galindo, C. and Fanning, C. M. 1998. The Pampean Orogeny of the southern proto-Andes evidence for Cambrian continental collision in the Sierras de Córdoba. En: Pankhurst, R. J. y Rapela, C. W. (Eds.), “The Proto-Andean Margin of Gondwana”, Special Publication N° 142:181 – 217, Geological Society, London.
- Spalletti, L. A., 1993. An iron-bearing wave-dominated siliciclastic shelf: Facies analysis and paleogeographic implications (Silurian-Lower Devonian Sierra Grande Formation, Southern Argentina). *Geological Journal*, 28: 137-148.
- Spalletti, L. A., Cingolani, C. A. and Varela, R., 1991. Ambientes y procesos generadores de las sedimentitas portadoras de hierro en la plataforma siluro-eodevónica de la Patagonia, República Argentina. *Revista Museo de La Plata, nueva serie, sección Geología*, 10: 305-318.
- Tickyj, H., Llambías, E.J. and Sato, A.M., 1999. El basamento cristalino de la región sur-oriental de la provincia de La Pampa: Extensión austral del Orógeno Famatiniano de Sierras Pampeanas. 14° Congreso Geológico Argentino, Actas 1: 160-163.
- Tickyj H., Llambías E. J. and Melchor, R. N., 2002. Ordovician Rocks from La Pampa Province, Argentina. En Aceñolaza, F. G. (ed.) *Aspects of the Ordovician System in Argentina*, Instituto Superior de Correlación Geológica, Serie Correlación Geológica 16:257-266. Tucumán.
- Varela, R.; Dalla Salda, L.; Cingolani C. and Gómez, V. 1991. Estructura, petrología y geocronología del basamento de la región del Limay, provincias de Río Negro y Neuquén, Argentina. *Revista Geológica de Chile*, 18 (2):147-163.
- Varela, R., Cingolani, C., Sato, A., Dalla Salda, L., Brito Neves, B.B., Basei, M.A.S., Siga Jr., O. and Teixeira, W., 1997. Proterozoic and Paleozoic evolution of Atlantic area of North Patagonian Massif Argentine. *South American Symposium on Isotope Geology (Sao Paulo/Brazil, June 15–18, 1997). Extended Abstracts*, 326–329.
- Varela, R., Basei, M.A.S., Sato, A.M., Siga, O. Jr., Cingolani, C.A. and Sato, K., 1998. Edades isotópicas Rb/Sr y U/Pb en rocas de Mina Gonzalito y Arroyo Salado. *Macizo Norpatagónico Atlántico, Río Negro, Argentina. X Congreso Latinoamericano de Geología y VI. Congreso Nacional de Geología Económica*, Actas I, 71–76.
- Varela, R., Basei, M.A.S., Sato, A.M., Llambías, E.J. and Siga Jr., O., 2001. Edades isotópicas neopaleozoicas del Complejo Yaminué, Macizo Norpatagónico. *Resúmenes 2° Simposio Argentino Paleozoico Superior*, 31 pp., Trelew.
- Zanettini, J. C. M., 1981. La Formación Sierra Grande (provincia de Río Negro). *Revista de la Asociación Geológica Argentina*, 36 (2):160-179.